

Chemical Block Ltd., offer on tRNA

<http://www.chemblock.com>, orders@chemical-block.com

Transfer Ribonucleic Acid: Arginine specific

lyophilized, from Escherichia coli (MRE 600)

Arginine Acceptor Activity: approx. 1400 picomoles per A260 unit

Transfer Ribonucleic Acid: Glutamic acid specific

lyophilized, from Escherichia coli (MRE 600)

Glutamic Acceptor Activity: approx. 1200 picomoles per A260 unit

Transfer Ribonucleic Acid: Lysine specific

lyophilized, from Escherichia coli (MRE 600)

Lysine Acceptor Activity: approx. 1500 picomoles per A260 unit

Transfer Ribonucleic Acid: Methionine specific

lyophilized, from Escherichia coli (MRE 600)

Methionine Acceptor Activity: approx. 1100 picomoles per A260 unit

Transfer Ribonucleic Acid: N-Formyl-methionine specific

lyophilized, from Escherichia coli (MRE 600)

Methionine Acceptor Activity: approx. 1500 picomoles per A260 unit

Transfer Ribonucleic Acid: Tyrosine specific

lyophilized, from Escherichia coli (MRE 600)

Tyrosine Acceptor Activity: approx. 1000 picomoles per A260 unit

Transfer Ribonucleic Acid: Valine specific

lyophilized, from Escherichia coli (MRE 600)

Valine Acceptor Activity: approx. 1400 picomoles per A260 unit

Transfer Ribonucleic Acid: Phenylalanine specific

lyophilized, from Escherichia coli (MRE 600)

Phenylalanine Acceptor Activity: approx. 1000 picomoles per A260 unit
1 mg =20 units A260

Transfer Ribonucleic Acid: Phenylalanine specific

lyophilized, from Bakers Yeast

Phenylalanine Acceptor Activity: approx. 1500 picomoles per A260 unit
1 mg =20 units A260

Transfer Ribonucleic Acid from Bakers Yeast

lyophilized powder,

approx. 19 A260 units per mg

Prepared by a modification of the procedure:

Holley, BBRC, 1963, 10, p.186-188